

Bridging the Gulf: 'Gulf War Syndrome' - what we know and what we don't

Despite considerable efforts it is likely that we will never know the full story of what has become known, albeit erroneously, as the 'Gulf War Syndrome' (GWS). However, some 16 years since the 1991 Gulf War, 44% of all United States (US) Gulf War veterans receive some form of disability payments, costing \$1 billion annually, and over 10% of United Kingdom (UK) Gulf veterans are in receipt of war pensions. So what is GWS and what might be its causes?

The ground phase of 1991 Gulf War lasted just four days and was a resounding military and medical military success. Traditional causes of non-battle casualties such as heat stroke and dehydration were rare. However, shortly after the war ended, reports emerged, initially from the US, of previously fit veterans developing unusual illnesses and symptoms. Media organisations quickly fuelled the emerging concerns by reporting increasing numbers of veterans having children with birth defects. Subsequently the US, and then the UK, began to conduct formal epidemiological research.

Initial studies used data from US health registries which provided systematic clinical evaluations; the process was subsequently repeated in the UK. Analysis of pooled data from over 100,000 programme attendees did not suggest any unusual pattern of illness, instead medically unexplained symptoms and syndromes were the most common diagnoses.^{1,2} However because programme attendees are self selecting the data obtained is only of limited scientific value. Nonetheless, if service during the 1991 Gulf War was associated with either a novel disease process or a dramatic elevation of a recognised but previously rare condition, then this would have been detected. Neither has happened. Furthermore, although the media reported an increase in the mortality rate in Gulf War veterans, numerous comprehensive analyses of the US and UK cohorts have not confirmed this other than an increase in accidental death, (US and UK) or suicide (US only) as observed in the aftermath of other conflicts.^{3,4}

Epidemiological reports of increased rates of symptom reporting in a cohort of US Gulf veterans found that symptom-defined conditions including chronic fatigue syndrome, depression, and post traumatic stress disorder were all elevated.⁵ The first UK systematic epidemiological study was undertaken by King's College London. This random sample of over 4,000 UK Gulf veterans was compared to similar numbers of active duty personnel who had deployed to Bosnia in 1992, and a further military non-deployed group.⁶ We found that the Gulf group were twice as likely to report each and every one of the 50 physical symptoms enquired about. Furthermore, the Gulf cohort reported decreased

health perception, but physical functioning was only very slightly different and still above expected non-military norms. Hence, Gulf veterans experienced more symptoms, endorsed more conditions, felt worse, but were still physically functioning almost as well as those deployed to another busy and stressful operational theatre.⁶

Other US, UK, Australian and Canadian epidemiological samples show essentially the same findings. Gulf War veterans report two to three times the rates of common symptoms as their non-deployed colleagues and also have more negative health perception and poorer quality of life.⁷ Nearly every study also confirms that the general increase in symptoms is not a new cluster of unusual symptoms specifically linked to Gulf service, suggesting that although subjective health has been clearly impaired, there is no specific nor unique 'Gulf War Syndrome'. A distinct syndrome was reported by US epidemiologist Robert Haley, but in a small study of a single unit with a low response rate and no controls.⁸ Furthermore, Haley's group has reported both central and peripheral nerve damage in the same veterans, which they attribute to exposure to a combination of chemical weapons and/or pesticides. However, expert review panels have not been convinced by either the medical evidence or the suggestion of significant exposure to chemical weapons. Our epidemiological study, and an even larger US study, failed to find evidence of significant damage to the peripheral nervous system, making exposure to organophosphate pesticides an unlikely cause of ill health.⁹

Yet, although there is no denying this change in symptoms and quality of life, it is equally clear that there has been no increase in well defined physical outcomes. For example, there has been no increase in cancer. All that has been found is a US study reporting an increase in motor neuron disease or amyotrophic lateral sclerosis (ALS) as it is known in the US.¹⁰ Irrespective of this, whilst ALS is a devastating disease, it remains very rare in veteran populations, and cannot account for more than a tiny fraction of the observed increase in morbidity in Gulf veterans. What Gulf veterans are therefore experiencing is an increase in symptoms, but not disease.

The search for possible aetiological agents has examined a variety of sub groups of deployed personnel. However the Gulf War health effect appears to have affected deployed military groups relatively equally. For instance, there is no consistent evidence of differences in the reporting of symptoms between the Services, suggesting that any possible causative agent of GWS would have to have equally affected those who operate over



Dr Amy Iversen is a Clinical Lecturer in Psychological Medicine at the Institute of Psychiatry with a special interest in veterans' health. She is particularly interested in the development of psychosocial interventions which address the complex needs of Gulf War veterans.



Ms Rachael Morris is a fourth year medical student at Guy's, King's and St. Thomas's School of Medicine, in London. She has a keen interest in mental health research and is currently rewriting a web-based encyclopaedia entry on Gulf War Syndrome.



Dr Neil Greenberg is a Military Psychiatrist who is a Senior Lecturer at King's College London. He has a keen interest in organisational mental health and, as a military officer, has a keen interest in psychological health in military personnel.



Prof Simon Wessely is a Professor of Epidemiology and Liaison Psychiatry in London. He has a keen interest in the investigation of contentious topics, including Gulf War Syndrome. He is currently the Director of the King's Centre for Military Health Research.

Gulf veterans experienced more symptoms, endorsed more conditions, felt worse, but were still physically functioning almost as well as those deployed to another busy and stressful operational theatre

Correspondence to:
Dr Neil Greenberg,
Academic Centre for Defence
Mental Health,
King's College London,
Weston Education Centre,
Cutcombe Rd,
London, SE5 9RJ, UK.
E. sososanta@aol.com

